


Elementarteilchenphysik – größte Messgeräte für kleinste Teilchen



ATLAS
EXPERIMENT
<http://atlas.ch>


Experimentelle Teilchen-
und Astroteilchenphysik

Was ist Teilchenphysik?



Demokrit (ca. 400 v.Chr.):

„Alle Dinge bestehen aus kleinen, unzerstörbaren Teilchen und aus leerem Raum“





Fragen:

- Was ist Materie?
- Wie entsteht Materie?
- Was hält Materie zusammen?

„Erforschen, was die Welt im Innersten zusammenhält“

Das Standardmodell der Teilchenphysik

Drei Generationen
der Materie (Fermionen)




	I	II	III		
Masse →	2,3 MeV	1,275 GeV	173,07 GeV	0	125,9 GeV
Ladung →	$\frac{2}{3}$	$\frac{2}{3}$	$\frac{2}{3}$	0	0
Spin →	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1	0
Name →	u up	c charm	t top	γ Photon	H Higgs Boson
	4,8 MeV	95 MeV	4,18 GeV	0	
	$-\frac{1}{3}$	$-\frac{1}{3}$	$-\frac{1}{3}$	0	
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1	
Quarks	d down	s strange	b bottom	g Gluon	
	<2 eV	<0,19 MeV	<18,2 MeV	91,2 GeV	
	0	0	0	0	
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1	
	ν_e Elektron- Neutrino	ν_μ Myon- Neutrino	ν_τ Tau- Neutrino	Z^0 Z Boson	
	0,511 MeV	105,7 MeV	1,777 GeV	80,4 GeV	
	-1	-1	-1	± 1	
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1	
Leptonen	e Elektron	μ Myon	τ Tau	W^\pm W Boson	Eichbosonen

Vom Kleinsten zum Größten

Verständnis der Welt
bei kleinsten Abständen



10^{-18} m



Verständnis der Welt
bei kosmischen Abständen


10^{25} m

Entstehung und Entwicklung
des Universums



Warum hohe Energien?

- ◆ Um neue, schwere Teilchen zu erzeugen
- ◆ Um kleinste Strukturen sichtbar zu machen



$$E = mc^2!$$





Elektronen-
mikroskop



Teilchen-
beschleuniger

Teilchenphysik = Hochenergiephysik

CERN in Genf...




Suche nach bisher unentdeckten
Teilchen (z.B. das Higgs-Boson)

Präzisionsmessungen
(Selbstkonsistenz?)

Teilchendetektoren


Teilchen werden mit hoher Energie aufeinander geschossen.

Detektoren weisen Kollisionsprodukte nach.




Was macht der Teilchenphysiker/ die Teilchenphysikerin?

● Analyse von Daten




Viele Milliarden
beobachtete
Teilchen-
kollisionen

Datenanalyse
(Software)



Zugrunde liegender physikalischer Prozess






Entwicklung und Bau von Detektoren


Die großen Teilchendetektoren können nicht von einer Gruppe alleine aufgebaut werden.

→ Große internationale Kollaborationen
(~50 - einige 1000 Mitglieder)

Einzelne Institute sind verantwortlich für einzelne Detektorkomponenten.



...





- | | |
|----------------|-------------|
| Argentina | Netherlands |
| Armenia | Norway |
| Australia | Poland |
| Austria | Portugal |
| Azerbaijan | Romania |
| Belarus | Russia |
| Brazil | Serbia |
| Canada | Slovakia |
| China | Slovenia |
| Czech Republic | Spain |
| Denmark | Sweden |
| France | Switzerland |
| Georgia | Taiwan |
| Germany | Turkey |
| Greece | UK |
| Israel | USA |
| Italy | CERN |
| Japan | JINR |
| Morocco | |

ATLAS Collaboration








Computertechnik

Detektoren müssen riesige Datenmengen
Verarbeiten und speichern
(viele 1000 Terabytes/Jahr für LHC-Experimente)

- Große und schnelle Computernetzwerke
- Aufwendige Speicherung der Daten
- Datenkomprimierung notwendig



Standard Model of FUNDAMENTAL PARTICLES AND INTERACTIONS


The Standard Model summarizes the current knowledge in Particle Physics. It is the quantum theory that includes the theory of strong interactions (quantum chromodynamics or QCD) and the unified theory of weak and electromagnetic interactions (electroweak). Gravity is included on this chart because it is one of the fundamental interactions even though not part of the "Standard Model."

FERMIONS

matter constituents
spin = 1/2, 3/2, 5/2, ...

Leptons spin = 1/2		
Flavor	Mass GeV/c ²	Electric charge
ν_e electron neutrino	<1×10 ⁻¹⁸	0
e electron	0.000511	-1
ν_μ muon neutrino	<0.0002	0
μ muon	0.106	-1
ν_τ tau neutrino	<0.02	0
τ tau	1.7771	-1

Quarks spin = 1/2		
Flavor	Approx. Mass GeV/c ²	Electric charge
u up	0.003	2/3
d down	0.006	-1/3
c charm	1.3	2/3
s strange	0.1	-1/3
t top	175	2/3
b bottom	4.3	-1/3



If the protons and neutrons in this picture were 10 cm across, then the quarks and electrons would be less than 0.1 mm in size and the entire atom would be about 10 km across.

BOSONS

force carriers
spin = 0, 1, 2, ...

Unified Electroweak spin = 1		
Name	Mass GeV/c ²	Electric charge
γ photon	0	0
W ⁻	80.4	-1
W ⁺	80.4	+1
Z ⁰	91.187	0

Strong (color) spin = 1		
Name	Mass GeV/c ²	Electric charge
g gluon	0	0

Color Charge
Each quark carries one of three types of "strong charge," also called "color charge." These charges have nothing to do with the colors of visible light. There are eight possible types of color charge for gluons. Just as electrically-charged particles interact by exchanging photons, in strong interactions color-charged particles interact by exchanging gluons. Leptons, photons, and W and Z bosons have no strong interactions and hence no color charge.

Quarks Confined in Mesons and Baryons

One cannot isolate quarks and gluons; they are confined in color-neutral particles called **hadrons**. This confinement (binding) results from multiple exchanges of gluons among the color-charged constituents. As color-charged particles (quarks and gluons) move apart, the energy in the color force field between them increases. This energy eventually is converted into additional quark-antiquark pairs (see figure below). The quarks and antiquarks then combine into hadrons; these are the particles seen to emerge. Two types of hadrons have been observed in nature: **mesons** $q\bar{q}$ and **baryons** qqq .

Residual Strong Interaction

The strong binding of color-neutral protons and neutrons to form nuclei is due to residual strong interactions between their color-charged constituents. It is similar to the residual electrical interaction that binds electrically neutral atoms to form molecules. It can also be viewed as the exchange of mesons between the hadrons.

PROPERTIES OF THE INTERACTIONS

Baryons qqq and Antibaryons $\bar{q}\bar{q}\bar{q}$					
Baryons are fermionic hadrons. There are about 140 types of baryons.					
Symbol	Name	Quark content	Electric charge	Mass GeV/c ²	Spin
p	proton	uud	1	0.938	1/2
\bar{p}	anti-proton	$\bar{u}\bar{u}\bar{d}$	-1	0.938	1/2
n	neutron	udd	0	0.940	1/2
Λ	lambda	uds	0	1.116	1/2
Ω^-	omega	sss	-1	1.672	3/2

Property	Interaction	Weak	Electromagnetic	Strong	
	Acts on:	Flavor	Electric Charge	Fundamental	Residual
Particles experiencing:	Mass - Energy	Quarks, Leptons	Electrically charged	Quarks, Gluons	Hadrons
Particles mediating:	Graviton (not yet observed)	W ⁺ W ⁻ Z ⁰	γ	Gluons	Mesons
Strength relative to electromag. for two u quarks at:	10 ⁻⁴¹	0.8	1	25	Not applicable to quarks
for two u quarks at:	10 ⁻⁴¹	10 ⁻⁴	1	60	Not applicable to quarks
for two protons in nucleus	10 ⁻³⁶	10 ⁻⁷	1	Not applicable to hadrons	20




Mesons $q\bar{q}$					
Mesons are bosonic hadrons. There are about 140 types of mesons.					
Symbol	Name	Quark content	Electric charge	Mass GeV/c ²	Spin
π^+	pion	u \bar{d}	+1	0.140	0
K ⁻	kaon	s \bar{u}	-1	0.494	0
ρ^+	rho	u \bar{d}	+1	0.770	1
B ⁰	B-zero	d \bar{b}	0	5.279	0
η_c	eta-c	c \bar{c}	0	2.980	0

Matter and Antimatter

For every particle type there is a corresponding antiparticle type, denoted by a bar over the particle symbol (unless 1 or - charge is shown). Particle and antiparticle have identical mass and spin but opposite charges. Some electrically neutral bosons (e.g., Z⁰, η , and $\eta_c = c\bar{c}$, but not K⁰ = d \bar{s}) are their own antiparticles.

Figures

These diagrams are an artist's conception of physical processes. They are not exact and have no meaningful scale. Green shaded areas represent the cloud of gluons or the gluon field, and red lines the quark paths.



The Particle Adventure

Visit the award-winning web feature *The Particle Adventure* at <http://ParticleAdventure.org>

This chart has been made possible by the generous support of:

U.S. Department of Energy
U.S. National Science Foundation
Lawrence Berkeley National Laboratory
Stanford Linear Accelerator Center
American Physical Society, Division of Particles and Fields
BURLE INDUSTRIES, INC.

©2000 Contemporary Physics Education Project. CPEP is a non-profit organization of teachers, physicists, and educators. Send mail to: CPEP, MS 50-100, Lawrence Berkeley National Laboratory, Berkeley, CA, 94720. For information on charts, text materials, hands-on classroom activities, and workshops, see:

<http://CPEPweb.org>